



DAIRYINFO

W-S Feed & Supplies, Ltd.
1805 Sawmill Road
Conestogo, ON N0B 1N0
Canada
1.800.265.2203
www.wsfeeds.ca

Taking service to another level!

SOIL COMPACTION...

The need to conduct field operations in a timely and efficient manner has renewed concerns regarding the “productivity robbing” effects of soil compaction. Today’s producer often manages more acres, which can sometimes pressure field work to be done when soils are wet. These factors are problematic on dairy and other livestock operations, where frequent and extensive manure application is necessary.

Typically, soils are 50% solid and 50% pore space, with about half of the pore space filled with water. Soils are most easily compacted when the soil water content is at, or just above, its field capacity. Pressure from wheel traffic and tillage consolidate the soil, first reducing the number and size of larger pores by re-arranging soil aggregates. Heavy loads can destroy the soil structure itself. The resulting soil has a greater density and lower porosity. Pores, especially the larger ones, are important for water and air movement. The potential for compaction increases as the soil-clay content increases; however, sandy soils can also become compacted, especially if sand particles are of different sizes.

Signs and symptoms of compaction can be found by examining the response of the soil and crops to wheel traffic. Compacted soils have imperfect drainage, resulting in ponding and increased runoff. Where the structure is destroyed the soil will be massive and cloddy. A horizontal or plate-like structure can also develop in the upper soil layer. The loss of structure disrupts natural pores and channels that are important for water and air movement, and the resulting increase in soil strength that impedes root proliferation.

Compaction effects are also exhibited in the growth of the plant. Uneven height is common where one plant appears normal and the adjacent plant is stunted. The root system will be malformed such that horizontal development occurs at the restrictive layer. Nutrient deficiencies, especially K, can develop in response to poorer soil aeration.

Compaction almost always causes a loss in yield. The magnitude of the yield loss is often related to the incidence of water stress conditions during the growing season.

A common assessment device is the penetrometer, a cone-tipped rod attached to a gauge that is pushed into the soil at a constant rate. It measures the resistance to penetration and somewhat simulates the environment that a growing root would experience. Simple penetrometers are pushed by hand and have a dial that translates the force into green, yellow, and red zones. Advanced units are mechanically driven and have sensors that can be calibrated to measure and record the resistance in units of pressure. The soil water content will have a significant impact on the penetration resistance. Soil bulk density is a useful measurement of compaction and its measurement provides for the calculation of porosity.

There is no critical level of any soil measurement that universally identifies the degree of soil compaction that will result in a crop yield reduction. Penetration resistance, as previously indicated, is greatly affected by soil water content and will be very high if readings are taken in dry conditions. Bulk density is not affected by moisture, but is a function of soil texture. It is recommended that any measurement should be made to develop a relative comparison between areas where compaction is suspected and where it is unlikely. For example, compare a headland with an area in the main part of the field, or a wheel-tracked area with a non-tracked area. It is often useful to excavate the soil to examine the soil structure and evaluate plant root distribution. Be sure to note the depth at which compaction occurs to determine the depth of the restrictive layer. Knowledge of the location and size of this layer will determine if deep tillage can remove the compaction. Because compaction has such a profound influence on soil tilth, its effects vary from year-to-year depending on weather. Clearly, yields are reduced in a dry year and similar in more normal years, if the soil isn’t compacted. However, yields are drastically reduced in compacted plots.

Whenever possible soil compaction should be avoided! Practices such as limiting operations on wet soils, reducing load weight when possible, and controlling traffic will go a long way toward limiting compaction and maintaining soil productivity. Often deep tillage or subsoiling is considered when compaction problems are severe. Some producers routinely subsoil as a form of primary tillage. Subsoiling can be conducted with a variety of tillage tools that will have a variable effect depending on soil conditions, the depth of tillage, and the tool used. Before deciding to subsoil it is important to diagnose the existence of compaction and to record the depth of the restrictive layer. If subsoiling is done it should be conducted 1-2 inches below the layer.

Soil compaction problems will continue being an issue in modern agriculture. Using common sense can help to avoid the occurrence of compaction: reduce loads; stay off wet soils; and control traffic. Maintaining soil fertility, especially with respect to K, is also important. Use a complete starter fertilizer for corn and be sure to re-supply crop K. Look for compaction symptoms and physically identify the existence of a restrictive layer before conducting subsoiling operations. Repeated use of a soil enhancement product can also prove to be very beneficial not only for improvement from compaction, but also for soil health in general. Do not abuse the soil in the fall, expecting that over-winter conditions will likely correct compaction.

(Edited from an article by R. Wolkowski ~ UW Extension)

Interested in discussing topics in this newsletter, or want to do a better job feeding and managing your cows? Call us!

Our goal is to help you. That's the W-S Feed commitment!

VOLUME 6 – Number 8 – August 2016

CONSIDER FALL SEEDING!

Fall is ideal for seeding certain forage crops – for a cutting before winter and, with some varieties, a cutting again in the spring prior to planting corn. Certain species are ideal for this scenario and might help to make a difference in your forage inventories throughout this fall and the coming winter. We're pleased to offer seed products such as triticale hybrids, forage oats, alfalfa and other seeds for late summer and fall planting... for outstanding results on your farm. Consider seeding a new alfalfa field, with a wide range of available varieties to meet almost any need or situation. Some of the grass species are also excellent to seed this time of year (see article to right). From August until October there is a "window of opportunity" open to help boost forage inventories! Why wait? Get the facts. Consider seeding possibilities this summer and fall!

FALL SEEDING GRASSES... BEST TIME?

Late-summer/fall establishment of grass is often desirable. Many producers do not realize how much fall seeding can impact the yield of grasses in the next growing year. Research trials using six forage grasses planted at several late-summer dates over three years, produced some interesting results. Seeding dates were spaced approximately every 2 to 3 weeks from August 1 to November 1. Species included orchardgrass, smooth brome grass, timothy, reed canarygrass, perennial ryegrass, and tall fescue. All grasses were seeded by mid-to-late September and produced stands with visible plants ahead of a killing frost most years, usually surviving the winter. Later seedings did not produce visible plants until spring, if at all. Slow establishing species like reed canarygrass produced better stands when seeded by early September. Timothy tended to be the most variable with regard to seeding date and next-year yield. The most important finding is that *earlier seeding dates* (early- to mid-August) usually had more tillers per square foot, more tillers per plant, and higher dry matter yield the following season. Delaying late summer seeding beyond this timeframe generally resulted in about one (1) ton/acre less yield the next year. These studies clearly show that delaying grass seeding in the late summer or early fall not only increases the risk of establishment failure but also reduces yields the next year. With this in mind, we recommend seeding grasses as early as possible during the month of August.



W-S Feed & Supplies, Ltd.
1805 Sawmill Road
Conestogo, ON N0B 1N0
Canada

www.wsfeeds.ca

Taking service to another level



***SOIL COMPACTION...
FALL SEEDING GRASSES... BEST TIME?
CONSIDER FALL SEEDING!
AUGUST 2016
CHECK IT OUT!***